

## **Anti Secretin (Porcine) Serum**

Cat. No. Y031

Lot No. 80271013

**Description:** This antiserum was raised in a rabbit by immunization with a carrier free synthetic secretin (porcine) peptide. The product vial contains 50  $\mu$ L of the titled antiserum obtained by lyophilizing its 0.001 M phosphate buffer (pH 7.0, 0.5mL) solution. It can be used for immunoassay, immunohistochemistry or any other immunoreaction with secretin.

**Immunogen:** Synthetic secretin (porcine), carrier free

**Host:** Rabbit

**Amino Acid Sequence of Secretin (porcine)<sup>1)</sup>:** HSDGTFTSEL SRLRDSARLQ RLLQGLV-NH<sub>2</sub>

**Product Form:** Lyophilized unpurified serum

**Size:** 50  $\mu$ L

**Reconstitution:** Reconstitute the product with 0.5mL of 0.01M PBS (pH 7.0) to make a 10 fold diluted stock solution. If it is stored in a refrigerator, add moderate antiseptic to the solution (e.g. NaN<sub>3</sub> 0.1%).

**Storage:** The product will be stable for over one year if it be stored at -20°C to -80°C until opened. Upon reconstitution, the antiserum solution must be stored at 2°C to 8°C and used within one month. Repeated freezing-thawing should be avoided.

**Suggested Working Dilution Range:** 1:2,000-13,500 (final dilution ~1:54,000) for radioimmunoassay; 1: 1,000-4,000 for immunohistochemistry (frozen or paraffin sections). Optimal dilution should be determined by each laboratory for each application.

**Specificity** (based on radioimmunoassay): Secretin (porcine) 100%, secretin (4-27) (porcine) 60%, secretin (14-27) (porcine) 10%, secretin (18-27) (porcine) <0.01%, VIP (porcine) 0%, gastrin (human) 0%, glucagon 0%

**Positive Control** (immunohistochemistry): Porcine or rat duodenum

**Species Tested:** Human, porcine, rat<sup>2)</sup>

### **REFERENCES:**

- 1) M. Carlquist, H. Joernvall et al., Human secretin is not identical to the porcine/bovine hormone. IRCS Medical Science 13:217-218, 1985
- 2) N. Yanaihara, M. Kubota et al., Synthesis of phenolic group containing analogues of porcine secretin and their immunological properties. Journal of Medicinal Chemistry 20: 648-655, 1977

**FOR RESEARCH LABORATORY USE ONLY**  
DO NOT USE ORGANIC SOLVENTS FOR DISSOLVING ANTISERUM



Yanaihara Institute Inc.

2480-1 Awakura, Fujinomiya, Shizuoka, 418-0011 Japan

TEL: +81-544-22-2771

FAX: +81-544-22-2770

For more information, please visit our internet website: <http://www.yanaihara.co.jp> or contact us to: [ask@yanaihara.co.jp](mailto:ask@yanaihara.co.jp)